

JOURNAL OF ALGEBRA, 10, 278-297 (1968)

On Commutativity of a Semigroup which is a Semilattice of Commutative Semigroups

REIKICHI YOSHIDA

Department of Mathematics, Ritumeikan University, Kyoto, Japan

AND

MIYUKI YAMADA

*Department of Mathematics, Shimane University, Matsue, Japan**Communicated by D. Rees**Received July 10, 1967*

Let $P_1(G)$ and $P_2(G)$ be abstract properties pertaining to commutative semigroups G in the sense of Cohn [3]. $P_1(G)$ is said to be weaker than or equal to $P_2(G)$ and denoted by $P_1(G) \geq P_2(G)$ if and only if, for any commutative semigroup S , $P_1(G)$ is satisfied by S (i.e., $P_1(S)$ is true) whenever $P_2(G)$ is satisfied by S . If $P_1(G) \geq P_2(G)$ and $P_2(G) \geq P_1(G)$, then $P_1(G)$ and $P_2(G)$ are said to be equivalent and denoted by $P_1(G) \equiv P_2(G)$. If $P_1(G) \equiv P_2(G)$, we regard $P_1(G)$ and $P_2(G)$ as the same property. When S is a semigroup which is a semilattice of commutative semi-groups S_ξ , $\xi \in \chi$, S is not necessarily commutative. However, there is an abstract property $P(G)$ pertaining to commutative semigroups G , such that, any semigroup which is a semilattice of commutative semigroups with $P(G)$ is commutative. Such an abstract property $P(G)$ is called a fully c-invariant property (abbrev., f.c.i.-property). For example, it is well-known (e.g., see Clifford [1]) that the property $P(G)$, " G is a group", is an f.c.i.-property. There is no greatest (i.e., weakest) f.c.i.-property with respect to the ordering relation defined above, but there is a maximal f.c.i.-property. Further, a maximal f.c.i.-property is not unique. The main purpose of this paper is to obtain maximal f.c.i.-properties, and some relevant results.

1. INTRODUCTION

A commutative idempotent semigroup Γ is called a *semilattice*. Define an ordering relation on Γ as follows:

$$(1.1) \quad \alpha \leq \beta \text{ if and only if } \alpha\beta = \beta\alpha = \beta.$$

Then, it is obvious that Γ is a partially ordered set with respect to \leq . If $\alpha \leq \beta$ and $\alpha \neq \beta$, then we shall denote it by $\alpha < \beta$. If Γ is a totally ordered set with respect to \leq , then Γ is called a *chain*. Now, let $\{S_\gamma : \gamma \in \Gamma\}$ (Γ : a semilattice) be a collection of semigroups S_γ . Then, each S_γ is called the γ -*component* of this collection. If γ is not a minimal element of Γ (i.e., if there is an element $\alpha \in \Gamma$ such that $\alpha < \gamma$), then the corresponding S_γ is called a *multiple-component*. Let $S = \sum\{S_\gamma : \gamma \in \Gamma\}$ (hereafter, \sum and $\dot{+}$ denote disjoint sum). If \circ is multiplication in S such that

(1.2) $S(\circ)$ is a semigroup, and each $S_\gamma (\gamma \in \Gamma)$ is embedded in $S(\circ)$, i.e., $x \circ y = xy$ for all $x, y \in S_\gamma$, and

$$(1.3) \quad S_\alpha \circ S_\beta \subset S_{\alpha\beta} \text{ for all } \alpha, \beta \in \Gamma,$$

then the resulting system $S(\circ)$ is called a *composition of $\{S_\gamma : \gamma \in \Gamma\}$ (with respect to Γ)*. Further, next we shall generalize this concept as follows: Let $\{S_\xi : \xi \in \chi\}$ (χ : a set) be a collection of semigroups S_ξ . Define multiplication $*$ in χ and multiplication \circ in $S = \sum\{S_\xi : \xi \in \chi\}$ such that $\chi(*)$ is a semilattice [chain] and $S(\circ)$ is a composition of $\{S_\xi : \xi \in \chi(*)\}$. In this case, $S(\circ)$ is called a *semilattice [linear] composition of $\{S_\xi : \xi \in \chi\}$* . Let $\{S_\gamma : \gamma \in \Gamma\}$ (Γ : a semilattice) be a collection of commutative semigroups S_γ . Then, sometimes there exists a composition $S(\circ)$ of $\{S_\gamma : \gamma \in \Gamma\}$ which is commutative. In this case, we shall call $S(\circ)$ a *commutative composition of $\{S_\gamma : \gamma \in \Gamma\}$* . Similarly if a semilattice [linear] composition $S(\circ)$ of a collection $\{S_\xi : \xi \in \chi\}$ (χ : a set) of commutative semigroups S_ξ is commutative, then $S(\circ)$ is called a *commutative semilattice [linear] composition of $\{S_\xi : \xi \in \chi\}$* . In general, for a given collection $\{S_\gamma : \gamma \in \Gamma\}$ (Γ : a semilattice) of semigroups S_γ there is not necessarily a composition of $\{S_\gamma : \gamma \in \Gamma\}$ (see [4]). If there exists at least one composition of $\{S_\gamma : \gamma \in \Gamma\}$, then the collection $\{S_\gamma : \gamma \in \Gamma\}$ is said to be *composable*. If Γ is a chain, then it is well-known (e.g., see [1]) that $\{S_\gamma : \gamma \in \Gamma\}$ is necessarily composable. For any given collection $\{S_\gamma : \gamma \in \Gamma\}$ (Γ : a semilattice) of commutative semigroups S_γ , a composition of $\{S_\gamma : \gamma \in \Gamma\}$ is (even if it exists) not necessarily commutative. This can be seen from the following simple example:

Let $\Gamma = \{\alpha, \beta\}$ ($\alpha\beta = \beta\alpha = \beta, \alpha \neq \beta$) be a chain, S_α a commutative semigroup, and S_β a null semigroup containing at least two elements. Let $S = S_\alpha \dot{+} S_\beta$, and define multiplication \circ in S as follows:

$$x \circ y = \begin{cases} xy & \text{if } x, y \in S_\alpha \text{ or } \in S_\beta, \\ y & \text{if } x \in S_\alpha, y \in S_\beta, \\ 0 & \text{if } x \in S_\beta, y \in S_\alpha, \end{cases}$$

where 0 is the zero element of S_β . Then $S(\circ)$ is a non-commutative composition of $\{S_\alpha, S_\beta\}$ with respect to Γ . In Section 2, we shall give a necessary and sufficient condition for a collection $\{S_\gamma : \gamma \in \Gamma\}$ (Γ : a semilattice)

of commutative semigroups S_γ to be composable. Further, in the case where $\{S_\gamma : \gamma \in \Gamma\}$ is composable, we shall give a method of construction of all compositions of $\{S_\gamma : \gamma \in \Gamma\}$. We also give a necessary and sufficient condition for $\{S_\gamma : \gamma \in \Gamma\}$ that every composition of $\{S_\gamma : \gamma \in \Gamma\}$ (if it exists) be necessarily commutative.

Let $P(G)$ be a proposition pertaining to commutative semigroups G . As in Cohn [3], $P(G)$ is said to be an *abstract property* (pertaining to commutative semigroups) if, and only if, $P(G)$ is invariant under isomorphism, i.e.,

(1.4) for any commutative semigroups S_1, S_2 such that $S_1 \cong S_2$ (S_1 is isomorphic with S_2), $P(S_1)$ is true whenever $P(S_2)$ is true and vice-versa.

If $P(S)$ is true for a commutative semigroup S , then we shall say that S *satisfies* $P(G)$. In this case, we also say that S is a commutative semigroup *with* $P(G)$. For example, the properties “ G is a group” and “ G is cancellative” (pertaining to commutative semigroups G) are abstract properties. Let $P_1(G)$ and $P_2(G)$ be abstract properties. Then $P_1(G)$ and $P_2(G)$ are said to be *equivalent* (denoted by $P_1(G) \equiv P_2(G)$) if the following (1.5) is fulfilled.

(1.5) For any commutative semigroup S , $P_1(S)$ is true if, and only if, $P_2(S)$ is true.

Hereafter, we shall consider $P_1(G)$ and $P_2(G)$ as the same property if they are equivalent. Define an ordering relation on the set \mathfrak{P} of abstract properties as follows: Let $P_1(G)$ and $P_2(G)$ be abstract properties. $P_1(G) \leq P_2(G)$ if the following (1.6) is fulfilled:

(1.6) For every commutative semigroup S , $P_2(S)$ is true whenever $P_1(S)$ is true.

If $P_1(G) \leq P_2(G)$ and $P_1(G) \not\equiv P_2(G)$, then the property $P_2(G)$ is said to be *weaker* than the property $P_1(G)$ and denoted by $P_1(G) < P_2(G)$. It is obvious that \mathfrak{P} is a partially ordered set with respect to this relation \leq (when we regard properties $P_1(G)$ and $P_2(G)$ as the same property if $P_1(G) \equiv P_2(G)$).

Next, consider the following propositions concerning an abstract property $P(G)$:

(1.7) For any collection $\{S_\gamma : \gamma \in \Gamma\}$ (Γ : a chain) of commutative semigroups S_γ , where each multiple-component S_α satisfies $P(G)$, every composition of $\{S_\gamma : \gamma \in \Gamma\}$ is commutative.

(1.8) For any collection $\{S_\gamma : \gamma \in \Gamma\}$ (Γ : a semilattice) of commutative semigroups S_γ , where each multiple-component S_α satisfies $P(G)$, every composition of $\{S_\gamma : \gamma \in \Gamma\}$ (if it exists) is commutative.

If (1.7) or (1.8) is true for $P(G)$, then $P(G)$ is called a *linearly c-extensible property* (abbrev., l.c.e.-property) or *fully c-extensible property* (abbrev.,

f.c.e.-property) respectively. For example, the abstract property "G is a group" pertaining to commutative semigroup G is an f.c.e.-property. By the definitions an f.c.e.-property is clearly an l.c.e.-property, but the converse is not true (see Remark below). In Section 3, we shall prove the existence of the weakest l.c.e.-property and the weakest f.c.e.-property and try to determine these properties.

Next, consider also the following propositions concerning an abstract property $P(G)$:

(1.9) For any collection $\{S_\xi : \xi \in \chi\}$ (χ : a set) of commutative semigroups S_ξ , where each S_ξ satisfies $P(G)$, every linear composition of $\{S_\xi : \xi \in \chi\}$ is commutative.

(1.10) For any collection $\{S_\xi : \xi \in \chi\}$ (χ : a set) of commutative semigroups S_ξ , where each S_ξ satisfies $P(G)$, every semilattice composition of $\{S_\xi : \xi \in \chi\}$ is commutative.

If (1.9) or (1.10) is true for $P(G)$, then $P(G)$ is called a *linearly c-invariant property* (abbrev., l.c.i.-property) or a *fully c-invariant property* (abbrev., f.c.i.-property) respectively. It is obvious from the definitions that an l.c.e. [f.c.e.]-property is an l.c.i. [f.c.i.]-property. In Section 4, we shall prove the existence of maximal l.c.i.-properties and maximal f.c.i.-properties and determine some of them.

Remark. Let $P_u(G)$ be an abstract property as follows:

(1.11) G is universal, i.e., $G^2 = G$.

Then it is easy to see that *universality* $P_u(G)$ is an l.c.e.-property (this will be shown later). Now, let T be a universal commutative semigroup which has a zero element 0 and whose annihilator A contains a non-zero element. The existence of such a semigroup T is easily seen from the following simple example: Let T^* be the semigroup consisting of all real numbers $\alpha > 1$ with respect to the usual multiplication. Then $I = \{\beta : \beta \geq 25, \beta \in T^*\}$ is an ideal of T^* . Let $T_1 = T^*/I$ be the Rees factor semigroup of T^* modulo I : $T_1 = (1, 25) \dot{+} \{O\}$, where O is the zero element of T_1 . Let $\Omega = \{(x, y) : x, y \in T_1, xy = O \text{ in } T_1\}$, and let $\Gamma = \Omega \setminus \{(x, y) : x, y \in (1, 25), x \times y = 25\}$ where $x \times y$ denotes the usual product of the real numbers x and y . Let $N = \{u, 0\}$ be a null semigroup of order 2, where 0 is the zero element of N . Let $T = N \dot{+} T_1 \setminus O$, and define multiplication \circ in T as follows: For $x, y \in T$,

$$x \circ y = \begin{cases} xy & \text{if } x, y \in T_1 \setminus O \text{ and } xy \neq O \text{ or if } x, y \in N, \\ 0 & \text{if } x \in N \text{ or } y \in N, \\ 0 & \text{if } x, y \in T_1 \setminus O, xy = O \text{ and } (x, y) \in \Gamma, \\ u & \text{if } x, y \in T_1 \setminus O, xy = O \text{ and } (x, y) \notin \Gamma. \end{cases}$$

Then T is a universal commutative semigroup which has a zero element, and whose annihilator contains a non-zero element.

Now, let $L_3 = \{\alpha, \beta, \gamma\}$ be a semilattice such that $\alpha < \gamma, \beta < \gamma, \alpha \not\leq \beta$ and $\beta \leq \alpha$. Let S_α and S_β be infinite cyclic semigroups generated by a and b respectively: $S_\alpha = (a)$ and $S_\beta = (b)$. Let $S_\gamma = T$. Then $S = S_\alpha \dot{+} S_\beta \dot{+} S_\gamma$ becomes a non-commutative composition of $\{S_\alpha, S_\beta, S_\gamma\}$ with respect to L_3 by the multiplication \odot defined as follows:

$$x \odot y = \begin{cases} xy & \text{if } x, y \in S_\alpha, \in S_\beta \text{ or } \in S_\gamma, \\ u & \text{if } x = a \text{ and } y = b, \\ 0 & \text{otherwise,} \end{cases}$$

where 0 is the zero element of $T(=S_\gamma)$ and u is a fixed non-zero element contained in the annihilator of $T(=S_\gamma)$. Hence $P_u(G)$ is not an f.c.e.-property.

Throughout this paper, if $\{S_\xi : \xi \in \chi\}$ is a collection of commutative semigroups S_ξ , we shall denote elements of S_ξ by small letters a_ξ, b_ξ, c_ξ etc. having ξ as their subscripts.

2. COMPOSITION THEOREMS

Let $\Omega = \{S_\gamma : \gamma \in \Gamma\}$ (Γ : a semilattice) be a collection of commutative semigroups S_γ . For every pair (α, β) of $\alpha, \beta \in \Gamma$ with $\alpha \leq \beta$, let $\mathfrak{M}(\alpha, \beta)$ be the set of mappings of S_α into S_β . Let $C(\alpha, \beta) = \{S_\xi : \xi \in \Gamma, \alpha \xi = \beta\}$. Clearly $S_\beta \in C(\alpha, \beta)$. For every $S_\xi \in C(\alpha, \beta)$, let ψ_ξ, φ_ξ be (not necessarily distinct) two mappings of S_ξ into $\mathfrak{M}(\alpha, \beta)$. Put $\psi_\xi(a_\xi) = \tilde{a}^{(\alpha, \beta)}$ and $\varphi_\xi(a_\xi) = \tilde{a}^{(\alpha, \beta)}$. Let $\mathfrak{M}_L(\Omega) = \mathfrak{M}_L(S_\gamma : \gamma \in \Gamma) = \{\tilde{a}^{(\alpha, \beta)} : \alpha \leq \beta, \alpha, \beta \in \Gamma, a_\xi \in S_\xi, S_\xi \in C(\alpha, \beta)\}$, and $\mathfrak{M}_R(\Omega) = \mathfrak{M}_R(S_\gamma : \gamma \in \Gamma) = \{\tilde{b}^{(\alpha, \beta)} : \alpha \leq \beta, \alpha, \beta \in \Gamma, a_\xi \in S_\xi, S_\xi \in C(\alpha, \beta)\}$. If

$$(2.1) \quad \mathfrak{M}(\Omega) = \mathfrak{M}(S_\gamma : \gamma \in \Gamma) = \mathfrak{M}_L(\Omega) \dot{+} \mathfrak{M}_R(\Omega)$$

satisfies the following condition (C), then $\mathfrak{M}(\Omega)$ is called a *set of composite factors on Ω* :

$$(C) \quad \left\{ \begin{array}{l} (1) \quad \tilde{a}_\alpha^{(\beta, \alpha\beta)} \tilde{c}_\gamma^{(\alpha\beta, \alpha\beta\gamma)} = \tilde{c}_\gamma^{(\beta, \beta\gamma)} \tilde{a}_\alpha^{(\beta\gamma, \alpha\beta\gamma)}, \\ (2) \quad \tilde{a}_\alpha^{(\alpha, \alpha)} = \tilde{a}_\alpha^{(\alpha, \alpha)} = \text{the inner translation } \rho_{a_\alpha} \text{ on } S_\alpha \\ \quad \text{induced by } a_\alpha, \\ (3) \quad \tilde{a}_\alpha^{(\beta, \alpha\beta)} \text{ and } \tilde{b}_\beta^{(\alpha, \alpha\beta)} \text{ are conjugate to each other} \\ \quad \text{in the following sense: } \tilde{a}_\alpha^{(\beta, \alpha\beta)}(b_\beta) = \tilde{b}_\beta^{(\alpha, \alpha\beta)}(a_\alpha). \end{array} \right.$$

THEOREM 1. Let $\Omega = \{S_\gamma : \gamma \in \Gamma\}$ (Γ : a semilattice) be a collection of commutative semigroups S_γ .

(i) Ω is composable if and only if there exists a set of composite factors on Ω .

(ii) Let $\mathfrak{M}(\Omega)$ of (2.1) be a set of composite factors on Ω . Then $S = \Sigma\{S_\gamma : \gamma \in \Gamma\}$ becomes a composition $S(\odot)$ of Ω by the multiplication \odot defined by

$$(2.2) \quad a_\alpha \odot b_\beta = \bar{a}_\alpha^{(\beta, \alpha\beta)}(b_\beta) (= \bar{b}_\beta^{(\alpha, \alpha\beta)}(a_\alpha)).$$

Further, every possible composition of Ω is found in this fashion.

Proof. (i) follows from (ii) and the definition of composability. Hence, we prove only (ii). Now let $\mathfrak{M}(\Omega)$ of (2.1) be a set of composite factors on Ω . For $a_\alpha, b_\beta, c_\gamma \in S$, $(a_\alpha \odot b_\beta) \odot c_\gamma = \bar{a}_\alpha^{(\beta, \alpha\beta)}(b_\beta) \odot c_\gamma = \bar{a}_\alpha^{(\beta, \alpha\beta)} \bar{c}_\gamma^{(\alpha\beta, \alpha\beta\gamma)}(b_\beta) = \bar{c}_\gamma^{(\beta, \beta\gamma)} \bar{a}_\alpha^{(\beta\gamma, \alpha\beta\gamma)}(b_\beta) = \bar{a}_\alpha^{(\beta\gamma, \alpha\beta\gamma)}(\bar{c}_\gamma^{(\beta, \beta\gamma)}(b_\beta)) = \bar{a}_\alpha^{(\beta\gamma, \alpha\beta\gamma)}(b_\beta \odot c_\gamma) = a_\alpha \odot (b_\beta \odot c_\gamma)$. Since $a_\alpha \odot b_\beta \in S_{\alpha\beta}$ for all $a_\alpha \in S_\alpha, b_\beta \in S_\beta$ and since $a_\alpha \odot b_\alpha = a_\alpha b_\alpha$ for all $a_\alpha, b_\alpha \in S_\alpha$, $S(\odot)$ is a composition of Ω . Conversely, let $S(\odot) = \Sigma\{S_\gamma : \gamma \in \Gamma\}$ be a composition of Ω . For $\alpha, \beta, \xi \in \Gamma$ with $\alpha \leq \beta, \alpha\xi = \beta$ and for $a_\xi \in S_\xi$, define mappings $\bar{a}_\xi^{(\alpha, \beta)} : S_\alpha \rightarrow S_\beta$ and $\bar{a}_\xi^{(\alpha, \beta)} : S_\alpha \rightarrow S_\beta$ as follows: $\bar{a}_\xi^{(\alpha, \beta)}(b_\alpha) = a_\xi \odot b_\alpha$ and $\bar{a}_\xi^{(\alpha, \beta)}(b_\alpha) = b_\alpha \odot a_\xi$. Then it is easy to see that $\mathfrak{M}(\Omega) = \{\bar{a}_\xi^{(\alpha, \beta)} : \alpha \leq \beta, a_\xi \in S_\xi \text{ with } \alpha\xi = \beta\} \cup \{\bar{a}_\xi^{(\alpha, \beta)} : \alpha \leq \beta, a_\xi \in S_\xi \text{ with } \alpha\xi = \beta\}$ satisfies (1)–(3) of the condition (C) in (2.1) and hence is a set of composite factors on Ω . Let $S(\circ)$ be the composition of Ω determined by this $\mathfrak{M}(\Omega)$ and (2.2). Then, $a_\alpha \odot b_\beta = \bar{a}_\alpha^{(\beta, \alpha\beta)}(b_\beta) = a_\alpha \odot b_\beta$. Hence $S(\odot) = S(\circ)$.

The composition $S(\circ)$ in (ii) of Theorem 1 is called the composition of Ω induced by $\mathfrak{M}(\Omega)$.

COROLLARY 1. Let $\Omega = \{S_\gamma : \gamma \in \Gamma\}$ (Γ : a semilattice) be a collection of commutative semigroups S_γ , and $\mathfrak{M}(\Omega)$ of (2.1) a set of composite factors on Ω . Then, the composition $S(\odot)$ of Ω induced by $\mathfrak{M}(\Omega)$ is non-commutative if and only if the following condition is satisfied:

$$(2.3) \quad \bar{a}_\alpha^{(\beta, \alpha\beta)} \neq \bar{a}_\alpha^{(\alpha\beta, \alpha\beta)} \text{ for some } a_\alpha \in S_\alpha, \alpha, \beta \in \Gamma.$$

Proof. If $\bar{a}_\alpha^{(\beta, \alpha\beta)} \neq \bar{a}_\alpha^{(\alpha\beta, \alpha\beta)}$, then there exists $b_\beta \in S_\beta$ such that $\bar{a}_\alpha^{(\beta, \alpha\beta)}(b_\beta) \neq \bar{a}_\alpha^{(\alpha\beta, \alpha\beta)}(b_\beta)$. Hence $a_\alpha \odot b_\beta \neq b_\beta \odot a_\alpha$ in $S(\odot)$. Conversely, suppose that the composition $S(\odot)$ of Ω induced by $\mathfrak{M}(\Omega)$ is non-commutative. Then $a_\xi \odot b_\eta \neq b_\eta \odot a_\xi$ for some $a_\xi \in S_\xi, b_\eta \in S_\eta, \xi, \eta \in \Gamma$. Hence $\bar{a}_\xi^{(\eta, \xi\eta)}(b_\eta) \neq \bar{a}_\xi^{(\eta, \xi\eta)}(b_\eta) = \bar{a}_\xi^{(\eta, \xi\eta)}(b_\eta)$. This implies that $\bar{a}_\xi^{(\eta, \xi\eta)} \neq \bar{a}_\xi^{(\eta, \xi\eta)}$.

COROLLARY 2. Let $\Omega = \{S_\gamma : \gamma \in \Gamma\}$ (Γ : a semilattice) be a collection of commutative semigroups S_γ . Then, every composition of Ω is commutative if and only if there is no set, $\mathfrak{M}(\Omega)$ of (2.1), of composite factors on Ω which satisfies the condition (2.3).

Now, as a special case, we consider a collection $\Omega = \{S_\gamma : \gamma \in \Gamma\}$ of commutative semigroups S_γ having a chain Γ as its index set. Let $\mathfrak{M}(\Omega)$ of (2.1) be a set of composite factors on Ω .

Then, we have the following lemmas:

LEMMA 1. For $\alpha, \beta \in \Gamma$ with $\alpha \leq \beta$, each of $\tilde{a}_\alpha^{(\beta, \alpha\beta)} (= \tilde{a}_\alpha^{(\beta, \beta)})$ and $\tilde{a}_\alpha^{(\beta, \alpha\beta)} (= \tilde{a}_\alpha^{(\beta, \beta)})$ is a translation on S_β .

Proof. It follows from (1) of (C) in (2.1) that $\tilde{a}_\alpha^{(\beta, \beta)}(b_\beta d_\beta) = \tilde{a}_\alpha^{(\beta, \beta)}(\tilde{d}_\beta^{(\beta, \beta)}(b_\beta)) = \tilde{d}_\beta^{(\beta, \beta)}\tilde{a}_\alpha^{(\beta, \beta)}(b_\beta) = \tilde{a}_\alpha^{(\beta, \beta)}\tilde{d}_\beta^{(\beta, \beta)}(b_\beta) = \tilde{d}_\beta^{(\beta, \beta)}(\tilde{a}_\alpha^{(\beta, \beta)}(b_\beta)) = \tilde{a}_\alpha^{(\beta, \beta)}(b_\beta) d_\beta$. Hence, $\tilde{a}_\alpha^{(\beta, \alpha\beta)} (= \tilde{a}_\alpha^{(\beta, \beta)})$ is a translation on S_β . Similarly, it can be easily proved that $\tilde{a}_\alpha^{(\beta, \alpha\beta)} (= \tilde{a}_\alpha^{(\beta, \beta)})$ is also a translation on S_β .

LEMMA 2. For $\alpha, \beta \in \Gamma$ with $\alpha \geq \beta$, $\tilde{a}_\alpha^{(\beta, \alpha)}(b_\beta) = \tilde{b}_\beta^{(\alpha, \alpha)}(a_\alpha)$ and $\tilde{b}_\alpha^{(\beta, \alpha)}(a_\beta) = \tilde{a}_\beta^{(\alpha, \alpha)}(b_\alpha)$.

Proof. This follows from (3) of (C) in (2.1).

Now, put $\tilde{a}_\alpha^{(\beta, \beta)} = \rho_{a_\alpha, \beta}$ and $\tilde{a}_\alpha^{(\beta, \beta)} = \sigma_{a_\alpha, \beta}$ for $\alpha \leq \beta$. Then

LEMMA 3. $\rho_{a_\alpha, \alpha} = \sigma_{a_\alpha, \alpha} =$ the inner translation ρ_{a_α} on S_α induced by a_α .

Proof. Obvious.

LEMMA 4. $\rho_{a_\alpha, \gamma}\sigma_{b_\beta, \gamma} = \sigma_{b_\beta, \gamma}\rho_{a_\alpha, \gamma}$ if $\alpha \leq \gamma, \beta \leq \gamma$.

Proof. $\rho_{a_\alpha, \gamma}\sigma_{b_\beta, \gamma} = \tilde{a}_\alpha^{(\gamma, \gamma)}\tilde{b}_\beta^{(\gamma, \gamma)} = \tilde{b}_\beta^{(\gamma, \gamma)}\tilde{a}_\alpha^{(\gamma, \gamma)}$ (by (1) of (C) in (2.1))
 $= \sigma_{b_\beta, \gamma}\rho_{a_\alpha, \gamma}$.

LEMMA 5.

$$\rho_{b_\beta}, \rho_{a_\alpha, \gamma} = \begin{cases} \rho_{\rho_{a_\alpha, \beta}(b_\beta), \gamma} & \text{if } \alpha \leq \beta \leq \gamma, \\ \rho_{\sigma_{b_\beta, \alpha}(a_\alpha), \gamma} & \text{if } \beta \leq \alpha \leq \gamma. \end{cases}$$

Proof. $\rho_{b_\beta, \gamma}\rho_{a_\alpha, \gamma} = \tilde{b}_\beta^{(\gamma, \gamma)}\tilde{a}_\alpha^{(\gamma, \gamma)}$. (The case $\alpha \leq \beta \leq \gamma$) $\rho_{b_\beta, \gamma}\rho_{a_\alpha, \gamma}(c_\gamma) = \tilde{b}_\beta^{(\gamma, \gamma)}\tilde{a}_\alpha^{(\gamma, \gamma)}(c_\gamma) = \tilde{a}_\alpha^{(\gamma, \gamma)}(\tilde{b}_\beta^{(\gamma, \gamma)}(c_\gamma)) = \tilde{a}_\alpha^{(\gamma, \gamma)}(\tilde{c}_\gamma^{(\beta, \gamma)}(b_\beta))$ (by Lemma 2) $= \tilde{c}_\gamma^{(\beta, \gamma)}\tilde{a}_\alpha^{(\gamma, \gamma)}(b_\beta) = \tilde{a}_\alpha^{(\beta, \beta)}\tilde{c}_\gamma^{(\beta, \gamma)}(b_\beta)$ (by (1) of (C) in (2.1)) $= \tilde{c}_\gamma^{(\beta, \gamma)}(\tilde{a}_\alpha^{(\beta, \beta)}(b_\beta)) = \tilde{a}_\alpha^{(\beta, \beta)}(b_\beta)^{(\gamma, \gamma)}(c_\gamma) = \rho_{\rho_{a_\alpha, \beta}(b_\beta), \gamma}(c_\gamma)$. Hence, $\rho_{b_\beta, \gamma}\rho_{a_\alpha, \gamma} = \rho_{\rho_{a_\alpha, \beta}(b_\beta), \gamma}$.

(The case $\beta \leq \alpha \leq \gamma$) $\rho_{b_\beta, \gamma}\rho_{a_\alpha, \gamma}(c_\gamma) = \tilde{b}_\beta^{(\gamma, \gamma)}\tilde{a}_\alpha^{(\gamma, \gamma)}(c_\gamma) = \tilde{a}_\alpha^{(\gamma, \gamma)}(\tilde{b}_\beta^{(\gamma, \gamma)}(c_\gamma)) = \tilde{a}_\alpha^{(\gamma, \gamma)}(\tilde{c}_\gamma^{(\beta, \gamma)}(b_\beta))$ (by Lemma 2) $= \tilde{c}_\gamma^{(\beta, \gamma)}\tilde{a}_\alpha^{(\gamma, \gamma)}(b_\beta) = \tilde{a}_\alpha^{(\beta, \alpha)}\tilde{c}_\gamma^{(\alpha, \gamma)}(b_\beta)$ (by (1) of (C) in (2.1)) $= \tilde{c}_\gamma^{(\alpha, \gamma)}(\tilde{a}_\alpha^{(\beta, \alpha)}(b_\beta)) = \tilde{c}_\gamma^{(\alpha, \gamma)}(\tilde{b}_\beta^{(\alpha, \alpha)}(a_\alpha))$ (by Lemma 2) $= \tilde{b}_\beta^{(\alpha, \alpha)}(a_\alpha)^{(\gamma, \gamma)}(c_\gamma)$ (by Lemma 2) $= \rho_{\tilde{b}_\beta^{(\alpha, \alpha)}(a_\alpha), \gamma}(c_\gamma) = \rho_{\sigma_{b_\beta, \alpha}(a_\alpha), \gamma}(c_\gamma)$.

Hence, $\rho_{b_\beta, \gamma}\rho_{a_\alpha, \gamma} = \rho_{\sigma_{b_\beta, \alpha}(a_\alpha), \gamma}$.

LEMMA 6.

$$\sigma_{b_\beta, \gamma} \sigma_{a_\alpha, \gamma} = \begin{cases} \sigma_{\sigma_{a_\alpha, \beta}(b_\beta), \gamma} & \text{if } \alpha \leq \beta \leq \gamma, \\ \sigma_{\rho_{b_\beta, \alpha}(a_\alpha), \gamma} & \text{if } \beta \leq \alpha \leq \gamma. \end{cases}$$

Proof. This can be proved by an analogous method to the proof of Lemma 5.

By Lemmas 3–6, we obtain the following result: Let $\Omega = \{S_\gamma : \gamma \in \Gamma\}$ (Γ : a chain) be a collection of commutative semigroups S_γ , $\mathfrak{M}(\Omega)$ of (2.1) a set of composite factors on Ω . Let $S(\circ)$ be the composition of $\{S_\gamma : \gamma \in \Gamma\}$ induced by $\mathfrak{M}(\Omega)$.

Then, there exists a system

(2.4) $\mathfrak{S}(\Omega) = \{\rho_{a_\alpha, \beta} : a_\alpha \in S_\alpha, \alpha \leq \beta, \alpha, \beta \in \Gamma\} \dot{+} \{\sigma_{a_\alpha, \beta} : a_\alpha \in S_\alpha, \alpha \leq \beta, \alpha, \beta \in \Gamma\}$, where $\rho_{a_\alpha, \beta}$ and $\sigma_{a_\alpha, \beta}$ are mappings of S_β into S_β , such that

$$(T) \begin{cases} (1) \rho_{a_\alpha, \beta} \text{ and } \sigma_{a_\alpha, \beta} \text{ are translations on } S_\beta, \\ (2) \rho_{a_\alpha, \alpha} = \sigma_{a_\alpha, \alpha} = \text{the inner translation } \rho_{a_\alpha} \text{ on } S_\alpha \\ \quad \text{induced by } a_\alpha, \\ (3) \rho_{a_\alpha, \gamma} \sigma_{b_\beta, \gamma} = \sigma_{b_\beta, \gamma} \rho_{a_\alpha, \gamma}, \\ (4) \rho_{b_\beta, \gamma} \rho_{a_\alpha, \gamma} = \begin{cases} \rho_{\rho_{a_\alpha, \beta}(b_\beta), \gamma} & \text{if } \alpha \leq \beta \leq \gamma, \\ \rho_{\sigma_{b_\beta, \alpha}(a_\alpha), \gamma} & \text{if } \beta \leq \alpha \leq \gamma, \end{cases} \\ (5) \sigma_{b_\beta, \gamma} \sigma_{a_\alpha, \gamma} = \begin{cases} \sigma_{\sigma_{a_\alpha, \beta}(b_\beta), \gamma} & \text{if } \alpha \leq \beta \leq \gamma, \\ \sigma_{\rho_{b_\beta, \alpha}(a_\alpha), \gamma} & \text{if } \beta \leq \alpha \leq \gamma. \end{cases} \end{cases}$$

Further, the multiplication \circ in $S(\circ)$ is represented by

$$(P) \quad a_\alpha \circ b_\beta = \begin{cases} \bar{a}_\alpha^{(\beta, \beta)}(b_\beta) = \rho_{a_\alpha, \beta}(b_\beta) & \text{if } \alpha \leq \beta, \\ \bar{b}_\beta^{(\alpha, \alpha)}(a_\alpha) = \sigma_{b_\beta, \alpha}(a_\alpha) & \text{if } \alpha \geq \beta. \end{cases}$$

In general, let $\Omega = \{S_\gamma : \gamma \in \Gamma\}$ (Γ : a chain) be a collection of commutative semigroups S_γ . For each pair (a_α, β) , where $a_\alpha \in S_\alpha$, $\alpha \leq \beta$ and $\alpha, \beta \in \Gamma$, let $\rho_{a_\alpha, \beta}$ and $\sigma_{a_\alpha, \beta}$ be (not necessarily distinct) two mappings of S_β into S_β . If $\mathfrak{S}(\Omega) = \{\rho_{a_\alpha, \beta} : a_\alpha \in S_\alpha, \alpha \leq \beta, \alpha, \beta \in \Gamma\} \dot{+} \{\sigma_{a_\alpha, \beta} : a_\alpha \in S_\alpha, \alpha \leq \beta, \alpha, \beta \in \Gamma\}$ satisfies (T) of (2.4), then $\mathfrak{S}(\Omega)$ is called a *factor set of translations on Ω* . From this definition and the above-mentioned result, we can conclude as follows: Let $\Omega = \{S_\gamma : \gamma \in \Gamma\}$ (Γ : a chain) be a collection of commutative semigroups S_γ . If $S(\circ) = \Sigma\{S_\gamma : \gamma \in \Gamma\}$ is a composition of $\Omega = \{S_\gamma : \gamma \in \Gamma\}$, then there exists a factor set of translations on Ω , say $\mathfrak{S}(\Omega) = \{\rho_{a_\alpha, \beta} : a_\alpha \in S_\alpha, \alpha \leq \beta, \alpha, \beta \in \Gamma\} \dot{+} \{\sigma_{a_\alpha, \beta} : a_\alpha \in S_\alpha, \alpha \leq \beta, \alpha, \beta \in \Gamma\}$, and \circ in $S(\circ)$ is represented by (P).

Conversely, let $\mathfrak{S}(\Omega) = \{\rho_{a_\alpha, \beta} : a_\alpha \in S_\alpha, \alpha \leq \beta, \alpha, \beta \in \Gamma\} \dot{+} \{\sigma_{a_\alpha, \beta} : a_\alpha \in S_\alpha, \alpha \leq \beta, \alpha, \beta \in \Gamma\}$ (Γ : a chain) be a factor set of translations on a collection $\Omega = \{S_\gamma : \gamma \in \Gamma\}$ of commutative semigroups S_γ . Next, we prove that

$S = \Sigma\{S_\gamma : \gamma \in \Gamma\}$ becomes a composition of Ω by the multiplication \odot given by (P). Let $S(\odot)$ be the set S in which multiplication \odot is defined by (P). $a_\alpha \odot b_\beta = \rho_{a_\alpha, \beta}(b_\beta) = a_\alpha b_\beta$. Since $S_\alpha \odot S_\beta \subset S_\beta$ if $\alpha \leq \beta$ and since $S_\alpha \odot S_\beta \subset S_\alpha$ if $\alpha \geq \beta$, in the both cases $S_\alpha \odot S_\beta \subset S_{\alpha\beta}$. Hence to prove $S(\odot)$ to be a composition of Ω , it is sufficient to prove the associativity of $S(\odot)$. Let α, β, γ be elements of Γ such that $\alpha \leq \beta \leq \gamma$.

(i) $(a_\alpha \odot b_\beta) \odot c_\gamma = \rho_{a_\alpha, \beta}(b_\beta) \odot c_\gamma = \rho_{\rho_{a_\alpha, \beta}(b_\beta), \gamma}(c_\gamma)$. On the other hand, $a_\alpha \odot (b_\beta \odot c_\gamma) = a_\alpha \odot (\rho_{b_\beta, \gamma}(c_\gamma)) = \rho_{a_\alpha, \gamma}(\rho_{b_\beta, \gamma}(c_\gamma)) = \rho_{b_\beta, \gamma}(\rho_{a_\alpha, \gamma}(c_\gamma))$. Thus by (4) of (T) in (2.4), $(a_\alpha \odot b_\beta) \odot c_\gamma = \rho_{\rho_{a_\alpha, \beta}(b_\beta), \gamma}(c_\gamma) = \rho_{b_\beta, \gamma}(\rho_{a_\alpha, \gamma}(c_\gamma)) = a_\alpha \odot (b_\beta \odot c_\gamma)$.

(ii) $(b_\beta \odot a_\alpha) \odot c_\gamma = \sigma_{a_\alpha, \beta}(b_\beta) \odot c_\gamma = \rho_{\sigma_{a_\alpha, \beta}(b_\beta), \gamma}(c_\gamma)$, while $b_\beta \odot (a_\alpha \odot c_\gamma) = b_\beta \odot (\rho_{a_\alpha, \gamma}(c_\gamma)) = \rho_{b_\beta, \gamma}(\rho_{a_\alpha, \gamma}(c_\gamma)) = \rho_{a_\alpha, \gamma}(\rho_{b_\beta, \gamma}(c_\gamma))$. Hence by (4) of (T) in (2.4), $(b_\beta \odot a_\alpha) \odot c_\gamma = \rho_{\sigma_{a_\alpha, \beta}(b_\beta), \gamma}(c_\gamma) = \rho_{a_\alpha, \gamma}(\rho_{b_\beta, \gamma}(c_\gamma)) = b_\beta \odot (a_\alpha \odot c_\gamma)$.

(iii) $(a_\alpha \odot c_\gamma) \odot b_\beta = \rho_{a_\alpha, \gamma}(c_\gamma) \odot b_\beta = \sigma_{b_\beta, \gamma}(\rho_{a_\alpha, \gamma}(c_\gamma)) = \rho_{a_\alpha, \gamma}(\sigma_{b_\beta, \gamma}(c_\gamma))$, while $a_\alpha \odot (c_\gamma \odot b_\beta) = a_\alpha \odot (\sigma_{b_\beta, \gamma}(c_\gamma)) = \rho_{a_\alpha, \gamma}(\sigma_{b_\beta, \gamma}(c_\gamma)) = \sigma_{b_\beta, \gamma}(\rho_{a_\alpha, \gamma}(c_\gamma))$. Hence by (3) of (T) in (2.4), $(a_\alpha \odot c_\gamma) \odot b_\beta = a_\alpha \odot (c_\gamma \odot b_\beta)$.

(iv) $(c_\gamma \odot a_\alpha) \odot b_\beta = \sigma_{a_\alpha, \gamma}(c_\gamma) \odot b_\beta = \sigma_{b_\beta, \gamma}(\sigma_{a_\alpha, \gamma}(c_\gamma)) = \sigma_{a_\alpha, \gamma}(\sigma_{b_\beta, \gamma}(c_\gamma))$, while $c_\gamma \odot (a_\alpha \odot b_\beta) = c_\gamma \odot (\rho_{a_\alpha, \beta}(b_\beta)) = \sigma_{\rho_{a_\alpha, \beta}(b_\beta), \gamma}(c_\gamma)$. Hence by (5) of (T) in (2.4), $(c_\gamma \odot a_\alpha) \odot b_\beta = c_\gamma \odot (a_\alpha \odot b_\beta)$.

(v) $(b_\beta \odot c_\gamma) \odot a_\alpha = \rho_{b_\beta, \gamma}(c_\gamma) \odot a_\alpha = \sigma_{a_\alpha, \gamma}(\rho_{b_\beta, \gamma}(c_\gamma)) = \rho_{b_\beta, \gamma}(\sigma_{a_\alpha, \gamma}(c_\gamma))$, while $b_\beta \odot (c_\gamma \odot a_\alpha) = b_\beta \odot (\sigma_{a_\alpha, \gamma}(c_\gamma)) = \rho_{b_\beta, \gamma}(\sigma_{a_\alpha, \gamma}(c_\gamma)) = \sigma_{a_\alpha, \gamma}(\rho_{b_\beta, \gamma}(c_\gamma))$. Hence by (3) of (T) in (2.4), $(b_\beta \odot c_\gamma) \odot a_\alpha = b_\beta \odot (c_\gamma \odot a_\alpha)$.

(vi) $(c_\gamma \odot b_\beta) \odot a_\alpha = \sigma_{b_\beta, \gamma}(c_\gamma) \odot a_\alpha = \sigma_{a_\alpha, \gamma}(\sigma_{b_\beta, \gamma}(c_\gamma)) = \sigma_{b_\beta, \gamma}(\sigma_{a_\alpha, \gamma}(c_\gamma))$, while $c_\gamma \odot (b_\beta \odot a_\alpha) = c_\gamma \odot (\sigma_{a_\alpha, \beta}(b_\beta)) = \sigma_{\sigma_{a_\alpha, \beta}(b_\beta), \gamma}(c_\gamma)$. Hence by (5) of (T) in (2.4), $(c_\gamma \odot b_\beta) \odot a_\alpha = c_\gamma \odot (b_\beta \odot a_\alpha)$.

In all cases, the associativity of the multiplication \odot in $S(\odot)$ can be proved.

Summerizing the results above, we obtain the following

THEOREM 2. *Let $\Omega = \{S_\gamma : \gamma \in \Gamma\}$ (Γ : a chain) be a collection of commutative semigroups S_γ . Let $\Xi(\Omega)$ of (2.4) be a factor set of translations on Ω . Then $S = \Sigma\{S_\gamma : \gamma \in \Gamma\}$ becomes a composition $S(\odot)$ of Ω by the multiplication \odot defined by (P). Further, every composition of Ω is found in this fashion.*

$S(\odot)$ in Theorem 2 is called the composition of Ω induced by $\Xi(\Omega)$. From Theorem 2, we obtain immediately the following

COROLLARY 1. *Let $\Omega = \{S_\gamma : \gamma \in \Gamma\}$ (Γ : a chain) be a collection of commutative semigroups S_γ , and $\Xi(\Omega)$ of (2.4) a factor set of translations on Ω . Then, the composition $S(\odot)$ of Ω induced by $\Xi(\Omega)$ is non-commutative if, and only if,*

$$(2.5) \quad \rho_{a_\alpha, \beta} \neq \sigma_{a_\alpha, \beta} \text{ for some } a_\alpha \in S_\alpha, \alpha, \beta \in \Gamma, \alpha < \beta.$$

Proof. Obvious.

COROLLARY 2. Let $\Omega = \{S_\gamma : \gamma \in \Gamma\}$ (Γ : a chain) be a collection of commutative semigroups S_γ . Every composition of Ω is commutative if and only if there is no factor set, $\Xi(\Omega)$ of (2.4), of translations on Ω which satisfies (2.5).

Proof. Obvious from Corollary 1.

In the case where every composition of a collection $\Omega = \{S_\gamma : \gamma \in \Gamma\}$ (Γ : a chain) of commutative semigroups S_γ is commutative, we have another construction theorem for the compositions of Ω which is somewhat simpler than Theorem 2:

THEOREM 3. Let $\Omega = \{S_\gamma : \gamma \in \Gamma\}$ (Γ : a chain) be a collection of commutative semigroups S_γ every composition of which is commutative. Let $S = \Sigma\{S_\gamma : \gamma \in \Gamma\}$. For each pair (α, β) , where $\alpha_\alpha \in S_\alpha$, $\alpha, \beta \in \Gamma$ and $\alpha \leq \beta$, let $\rho_{\alpha_\alpha, \beta}$ be a mapping of S_β into S_β . Let $\Xi(\Omega) = \{\rho_{\alpha_\alpha, \beta} : \alpha_\alpha \in S_\alpha, \alpha \leq \beta, \alpha, \beta \in \Gamma\}$. If $\Xi(\Omega)$ satisfies the condition

$$(\bar{T}) \begin{cases} (1) & \rho_{\alpha_\alpha, \beta} \text{ is a translation on } S_\beta, \\ (2) & \rho_{\alpha_\alpha, \alpha} = \text{the inner translation } \rho_{\alpha_\alpha} \text{ on } S_\alpha \text{ induced by } a_\alpha, \\ (3) & \rho_{\alpha_\alpha, \gamma} \rho_{b_\beta, \gamma} = \rho_{b_\beta, \gamma} \rho_{\alpha_\alpha, \gamma} = \rho_{\rho_{\alpha_\alpha, \beta}(b_\beta), \gamma} \text{ if } \alpha \leq \beta \leq \gamma, \end{cases}$$

then S becomes a composition $S(\odot)$ of Ω by the multiplication \odot defined by

$$(\bar{P}) \quad a_\alpha \odot b_\beta = b_\beta \odot a_\alpha = \rho_{a_\alpha, \beta}(b_\beta) \quad \text{if } \alpha \leq \beta.$$

Further, every composition of Ω is found in this fashion.

Proof. The first half of the theorem: Let $\sigma_{\alpha_\alpha, \beta} = \rho_{\alpha_\alpha, \beta}$, and put $\Xi(\Omega) = \{\rho_{\alpha_\alpha, \beta} : \alpha_\alpha \in S_\alpha, \alpha \leq \beta, \alpha, \beta \in \Gamma\} \dot{+} \{\sigma_{\alpha_\alpha, \beta} : \alpha_\alpha \in S_\alpha, \alpha \leq \beta, \alpha, \beta \in \Gamma\}$. Then $\Xi(\Omega)$ is clearly a factor set of translations on Ω . Hence, S becomes a composition of Ω by the multiplication \odot defined by (P). On the other hand, in this case it is easy to see that (P) means (\bar{P}) .

The second half of the theorem: Suppose that $S(\odot)$ is a composition of $\{S_\gamma : \gamma \in \Gamma\}$. Then, by the assumption, $S(\odot)$ is commutative. For any $a_\alpha \in S_\alpha$, $\alpha, \beta \in \Gamma$ with $\alpha \leq \beta$, define a mapping $\rho_{\alpha_\alpha, \beta} : S_\beta \rightarrow S_\beta$ by $\rho_{\alpha_\alpha, \beta}(b_\beta) = a_\alpha \odot b_\beta = b_\beta \odot a_\alpha$. Then it is easy to see that $\Xi(\Omega) = \{\rho_{\alpha_\alpha, \beta} : \alpha_\alpha \in S_\alpha, \alpha \leq \beta, \alpha, \beta \in \Gamma\}$ satisfies (\bar{T}) and the multiplication \odot in $S(\odot)$ is given by (\bar{P}) .

Next, we present some results concerning a factor set of translations on a collection $\Omega = \{S_\gamma : \gamma \in \Gamma\}$ (Γ : a chain) of commutative semigroups S_γ .

LEMMA 7. Let $\Omega = \{S_\gamma : \gamma \in \Gamma\}$ (Γ : a chain) be a collection of commutative semigroups S_γ . Let $\Xi(\Omega)$ of (2.4) be a factor set of translations on Ω which satisfies (T) in (2.4). Then,

- (i) for any $\alpha_\alpha \in S_\alpha$, $\alpha, \beta \in \Gamma$ with $\alpha \leq \beta$, $y_\beta \rho_{\alpha_\alpha, \beta}(x_\beta) = \sigma_{\alpha_\alpha, \beta}(y_\beta) x_\beta$, i.e., $\rho_{\alpha_\alpha, \beta}$ and $\sigma_{\alpha_\alpha, \beta}$ are linked, and
- (ii) $\rho_{\alpha_\alpha, \beta} \mid S_\beta^2 = \sigma_{\alpha_\alpha, \beta} \mid S_\beta^2$.¹

¹ $\rho_{\alpha_\alpha, \beta} \mid S_\beta^2$ denotes the restriction of $\rho_{\alpha_\alpha, \beta}$ to S_β^2 .

Proof. $y_\beta \rho_{a_\alpha, \beta}(x_\beta) = \rho_{y_\beta, \beta}(\rho_{a_\alpha, \beta}(x_\beta))$ (by (2) of (T)) $= \rho_{a_\alpha, \beta} \rho_{y_\beta, \beta}(x_\beta) = \rho_{\sigma_{a_\alpha, \beta}(y_\beta), \beta}(x_\beta)$ (by (4) of (T)) $= \sigma_{a_\alpha, \beta}(y_\beta) x_\beta$ (by (2) of (T)).

(ii) $\rho_{a_\alpha, \beta}(x_\beta y_\beta) = \rho_{a_\alpha, \beta}(x_\beta) y_\beta = y_\beta \rho_{a_\alpha, \beta}(x_\beta) = \sigma_{a_\alpha, \beta}(y_\beta) x_\beta$ (by (i)) $= x_\beta \sigma_{a_\alpha, \beta}(y_\beta) = \sigma_{a_\alpha, \beta}(x_\beta y_\beta)$. Hence $\rho_{a_\alpha, \beta} \mid S_\beta^2 = \sigma_{a_\alpha, \beta} \mid S_\beta^2$.

A commutative semigroup S is said to be *reductive* if it satisfies the following abstract property $P_r(G)$:

Reductivity $P_r(G)$: $ax = bx$ for all $x \in G$ implies $a = b$.

LEMMA 8. Let Ω and $\Xi(\Omega)$ be as in Lemma 7. If each of the multiple-components of Ω is universal or reductive, then $\rho_{a_\alpha, \beta} = \sigma_{a_\alpha, \beta}$ for any $a_\alpha \in S_\alpha$, $\alpha, \beta \in \Gamma$ with $\alpha \leq \beta$.

Proof. If $\alpha = \beta$, then clearly $\rho_{a_\alpha, \beta} = \sigma_{a_\alpha, \beta} = \rho_{a_\alpha}$. Suppose that $\alpha < \beta$. If S_β is universal, then $\rho_{a_\alpha, \beta} \mid S_\beta^2 = \rho_{a_\alpha, \beta} \mid S_\beta^2 = \sigma_{a_\alpha, \beta} \mid S_\beta^2 = \sigma_{a_\alpha, \beta}$. If S_β is reductive, then by Lemma 7 $\rho_{a_\alpha, \beta}(a_\beta b_\beta) = \sigma_{a_\alpha, \beta}(a_\beta b_\beta)$ for any a_β, b_β . Hence $\rho_{a_\alpha, \beta}(a_\beta) b_\beta = \sigma_{a_\alpha, \beta}(a_\beta) b_\beta$ for all $b_\beta \in S_\beta$. By the reductivity of S_β , $\rho_{a_\alpha, \beta}(a_\beta) = \sigma_{a_\alpha, \beta}(a_\beta)$ for all $a_\beta \in S_\beta$. This implies that $\rho_{a_\alpha, \beta} = \sigma_{a_\alpha, \beta}$.

By using Lemma 8 and Corollary 2 to Theorem 2, we obtain

COROLLARY. Let $\Omega = \{S_\gamma : \gamma \in \Gamma\}$ (Γ : a chain) be a collection of commutative semigroups S_γ . If each of the multiple-components of Ω is universal or reductive, then every composition of Ω is commutative.

Proof. Obvious.

Remark. This result will be more generalized in the next section.

3. THE WEAKEST F.C.E. [L.C.E.]-PROPERTY

In this section, we investigate l.c.e.-properties and f.c.e.-properties.

Let us consider the following abstract property $P_q(G)$ pertaining to commutative semigroups G :

(3.1) There is no system $\{\sigma, \rho\}$ of distinct two translations on G such that (1) $\sigma\rho = \rho\sigma$ and (2) $\sigma \mid G^2 = \rho \mid G^2$.

This property $P_q(G)$ is called *quasi-reductivity*. As is shown later, reductivity implies quasi-reductivity. However, the converse is not true.

LEMMA 9. Let $\{\sigma, \rho\}$ be a system of distinct two translations σ, ρ on a commutative semigroup S such that $\sigma\rho = \rho\sigma$ and $\sigma \mid S^2 = \rho \mid S^2$. Then there exist distinct two elements $x, y \in S$ and a prime element² $t \in S$ such that (1) $xa = ya$ for all $a \in S$ and (2) $\sigma(t) = x$ and $\rho(t) = y$.

² An element of $S \setminus S^2$ is called prime.

Proof. Since $\sigma \mid S^2 = \rho \mid S^2$ and $\sigma \neq \rho$, $\sigma(t) \neq \rho(t)$ for some $t \in S \setminus S^2$. It is obvious that t is a prime element of S . Putting $\sigma(t) = x$ and $\rho(t) = y$, we have $xa = \sigma(t)a = \sigma(ta) = \rho(t)a = ya$ for all $a \in S$.

Example. Let $S = \{a, a^2, \dots, a^n\}$ be a cyclic semigroup of order n such that $n \geq 2$, $a^{n-1} \neq a^n$ and $aa^n = a^n$. Define mappings $\rho, \sigma : S \rightarrow S$ as follows: $\rho(a) = a^{n-1}$, $\rho(a^i) = a^n$ if $i > 1$; and $\sigma(a^i) = a^n$ for all i . Then ρ, σ are translations on S such that $\sigma\rho = \rho\sigma$ and $\sigma \mid S^2 = \rho \mid S^2$. Hence, of course, S is not quasi-reductive.

By using Lemma 9, we can prove the following theorem:

THEOREM 4. $P_q(G)$ is the weakest l.c.e.-property.

Proof. At first, we prove that $P_q(G)$ is an l.c.e.-property. Suppose that $P_q(G)$ is not an l.c.e.-property. Then there exists a collection $\Omega = \{S_\gamma : \gamma \in \Gamma\}$ (Γ : a chain) of commutative semigroups S_γ such that $P_q(S_\alpha)$ is true for every multiple-component S_α of Ω and some composition $S(\circ)$ ($S = \Sigma\{S_\gamma : \gamma \in \Gamma\}$) of Ω is non-commutative. Hence, $a \circ b \neq b \circ a$ for some $a \in S_\gamma$, $b \in S_\beta$, $\gamma < \beta$. Now, define mappings $\rho, \sigma : S_\beta \rightarrow S_\beta$ as follows: $\rho(x) = a \circ x$ and $\sigma(x) = x \circ a$ for $x \in S_\beta$. Then ρ, σ are translations on S_β . For any $x \in S_\beta$, $\rho\sigma(x) = \sigma(\rho(x)) = \sigma(a \circ x) = (a \circ x) \circ a = a \circ (x \circ a) = a \circ \sigma(x) = \rho(\sigma(x)) = \sigma\rho(x)$. Thus $\rho\sigma = \sigma\rho$. Further for any $x, y \in S_\beta$, $\rho(xy) = a \circ xy = (a \circ x) \circ y = (a \circ x)y = y(a \circ x) = (y \circ a) \circ x = (y \circ a)x = x(y \circ a) = (x \circ y) \circ a = xy \circ a = \sigma(xy)$. Hence $\rho \mid S_\beta^2 = \sigma \mid S_\beta^2$. This contradicts to the assumption that S_β satisfies the property $P_q(G)$. Hence, $P_q(G)$ must be an l.c.e.-property. Next, suppose that there exists an l.c.e.-property $P(G)$ such that $P(G) \leq P_q(G)$. Then there exists a commutative semigroup S_0 such that $P(S_0)$ is true and $P_q(S_0)$ is not true. Since $P_q(S_0)$ is not true, there is a system $\{\sigma, \rho\}$ of distinct two translations σ, ρ on S_0 such that $\sigma\rho = \rho\sigma$ and $\sigma \mid S_0^2 = \rho \mid S_0^2$. Accordingly, there exist elements x, y, t of S_0 which satisfy (1), (2) of Lemma 9. Now, let $S_1 = (a)$ be an infinite cyclic semigroup generated by a and let $L = \{0, 1\}$ be a chain with respect to the usual multiplication. Define multiplication \circ in $S = S_1 \dot{+} S_0$ as follows: $a^i \circ u = \rho^i(u)$ for $u \in S_0$; $u \circ a^i = \sigma^i(u)$ for $u \in S_0$; $v \circ w = vw$ if $v, w \in S_1$ or $\in S_0$. Then $S(\circ)$ is a composition of $\{S_1, S_0\}$ with respect to L . Since $a \circ t = \rho(t) = y \neq x = \sigma(t) = t \circ a$, $S(\circ)$ is non-commutative. This contradicts to the assumption that $P(G)$ is an l.c.e.-property. Hence $P(G) \leq P_q(G)$.

COROLLARY. Each of reductivity, universality and the property "reductive or universal" is an l.c.e.-property.

Proof. In general, it is easy to see that if $P(G)$ and $P_1(G)$ are abstract properties such that $P_1(G) \leq P(G)$, and if $P(G)$ is an l.c.e.-property, then $P_1(G)$ is also an l.c.e.-property. For abstract properties $P_1(G)$ and $P_2(G)$,

denote the property " $P_1(G)$ or $P_2(G)$ " by $P_1(G) \vee P_2(G)$. It is obvious that $P_1(G) \leq P_1(G) \vee P_2(G)$ and $P_2(G) \leq P_1(G) \vee P_2(G)$. Now, we prove that $P_r(G) \vee P_u(G) \leq P_q(G)$. Suppose that $P_r(S) \vee P_u(S)$ is true for a commutative semigroup S . Since $P_r(S) \vee P_u(S)$ is true, at least one of $P_r(S)$ and $P_u(S)$ is true. If $P_q(S)$ is not true, then there is a system $\{\sigma, \rho\}$ of distinct two translations σ, ρ on S such that $\sigma\rho = \rho\sigma$ and $\sigma \mid S^2 = \rho \mid S^2$. Hence, by Lemma 9, there exist $x, y, t \in S$ which satisfy the conditions (1), (2) of Lemma 9. This contradicts to the assumption that S is reductive or universal. Hence, $P_q(S)$ must be true. Since $P_r(G) \leq P_r(G) \vee P_u(G) \leq P_q(G)$ and $P_u(G) \leq P_r(G) \vee P_u(G) \leq P_q(G)$ and since $P_q(G)$ is an l.c.e.-property, each of $P_r(G)$, $P_u(G)$ and $P_r(G) \vee P_u(G)$ is also an l.c.e.-property.

Remarks. (1) Moreover, the following is obvious from Theorem 4: Let $\Omega = \{S_\xi : \xi \in X\}$ (X : a set) be a collection of commutative semigroups S_ξ , where $P_q(S_\xi)$ is true for all $S_\xi \in \Omega$. Then, every linear composition of Ω is commutative.

(2) For a special collection $\Omega = \{S_\gamma : \gamma \in \Gamma\}$ (Γ : a chain) of commutative semigroups S_γ , every composition of Ω is commutative even if there exists a multiple-component S_λ which does not satisfy $P_q(G)$. For example, let $L = \{0, 1\}$ be a chain with respect to the usual multiplication, $S_1 = \{e\}$ a semigroup consisting of a single element e and $S_0 = \{a, a^2, \dots, a^{n-1}, a^n\}$ a cyclic semigroup of order n ($n > 2$) such that $a^{n-1} \neq a^n$ and $aa^n = a^n$. Then, it is easy to see from the above-mentioned example that $P_q(S_0)$ is not true. However, there is no non-commutative composition of $\{S_1, S_0\}$ with respect to L .

Hereafter, for any element x of a commutative semigroup S , the inner translation on S induced by x will be denoted by ρ_x .

Now, let us consider the following abstract property $P_r^*(G)$ pertaining to commutative semigroups G :

(3.2) There is no system $\{u, v; \xi, \eta\}$ of distinct two elements u, v of G and (not necessarily distinct) translations ξ, η on G such that (1) $\xi\eta = \eta\xi = \rho_u = \rho_v$ and (2) $\xi(u) = \xi(v)$ and $\eta(u) = \eta(v)$.

LEMMA 10. Let $\{u, v; \xi, \eta\}$ be a system of distinct two elements u, v of a commutative semigroup S and translations ξ, η on S , satisfying (1), (2) of (3.2). Then, $uz = vz$ for all $z \in S$.

Proof. Since $\xi\eta = \eta\xi = \rho_u = \rho_v$, it follows that $\rho_u(z) = \rho_v(z)$, i.e. $uz = vz$ for all $z \in S$.

LEMMA 11. $P_r^*(G)$ is equivalent to $P_r(G)$.

Proof. At first, we prove that $P_r(G) \leq P_r^*(G)$. Suppose that $P_r^*(S)$ is not

true for some commutative semigroup S . Then by Lemma 10, there exist distinct two elements u, v of S such that $uz = vz$ for all $z \in S$. Hence S is not reductive, and accordingly $P_r(S)$ is not true. Thus, $P_r(G) \leq P_r^*(G)$. Conversely, suppose that $P_r(S)$ is not true for some commutative semigroup S . Then, there exist $u, v \in S$ such that $uz = vz$ for all $z \in S$.

The case $S^2 \ni u$ or $\ni v$. Assume that $S^2 \ni u$, without the loss of generality. Let $u = cd$, $c, d \in S$, and let $\xi = \rho_c$ and $\eta = \rho_d$. Then $\xi\eta = \rho_c\rho_d = \rho_{cd} = \rho_u = \rho_v$ and $\eta\xi = \rho_d\rho_c = \rho_{dc} = \rho_{cd} = \rho_u = \rho_v$. Further, $\xi(u) = \rho_c(u) = uc = vc = \rho_c(v) = \xi(v)$ and $\eta(u) = \rho_d(u) = ud = vd = \rho_d(v) = \eta(v)$. Hence, there exists a system $\{u, v; \xi, \eta\}$ which satisfies (1), (2) of (3.2).

The case $S^2 \not\ni u, v$. Let $\xi = \rho_u$, and let η be a mapping: $S \rightarrow S$ such that $\eta(u) = v$, $\eta(v) = u$ and $\eta(x) = x$ if $x \neq u, v$. It is easy to see that η is a translation on S . (Note that $ux \neq u, v$ and $vx \neq v, u$ for all $x \in S$). At first, $\xi\eta = \rho_u$. In fact, $\rho_u\eta(u) = \eta(\rho_u(u)) = \eta(u^2) = uu = \rho_u(u)$, $\rho_u\eta(v) = \eta(\rho_u(v)) = \eta(uv) = uv = vu = \rho_u(v)$, and $\rho_u\eta(x) = \eta(\rho_u(x)) = \eta(xu) = xu = \rho_u(x)$ if $x \neq u, v$. Hence $\rho_u = \rho_u\eta = \xi\eta$. Next, $\eta\xi = \rho_u$. In fact, $\eta\rho_u(u) = \rho_u(\eta(u)) = \rho_u(v) = vu = uv = uu = \rho_u(u)$, $\eta\rho_u(v) = \rho_u(\eta(v)) = \rho_u(u) = \rho_u(v)$, and $\eta\rho_u(x) = \rho_u(\eta(x)) = \rho_u(x)$ if $x \neq u, v$. Hence $\rho_u = \eta\rho_u = \eta\xi$. Consequently, we have $\xi\eta = \rho_u = \rho_v = \eta\xi$. Further, $\xi(u) = \rho_u(u) = uu = vu = \rho_u(v) = \xi(v)$, and $\eta(u) = v = \eta(v)$. Thus, the system $\{u, v; \xi, \eta\}$ satisfies (1), (2) of (3.2). This means that $P_r^*(S)$ is not true. Hence, it follows that $P_r^*(G) \leq P_r(G)$.

Since $P_r^*(G) \leq P_r(G)$ and $P_r(G) \leq P_r^*(G)$, we have $P_r^*(G) = P_r(G)$.

By using Lemmas 10 and 11, we can prove the following theorem which is one of the main results of this paper:

THEOREM 5. $P_r(G)$ is the weakest f.c.e.-property.

Proof. At first, we prove that $P_r(G)$ is an f.c.e.-property. Let $\Omega = \{S_\gamma : \gamma \in \Gamma\}$ (Γ : a semilattice) be a collection of commutative semigroups S_γ such that every multiple-component S_ξ satisfies $P_r(G)$. Let $S(\odot)$ be any composition of Ω . For any $x, y \in S(\odot)$, we next prove that $x \odot y = y \odot x$. Let $x \in S_\alpha$ and $y \in S_\beta$.

Case 1. ($\alpha = \beta$). In this case, $x \odot y = xy = yx = y \odot x$.

Case 2. ($\alpha \neq \beta$). In this case, $P_r(S_{\alpha\beta})$ is true since $S_{\alpha\beta}$ is a multiple-component. For $a, b \in S_{\alpha\beta}$, we have $(x \odot y)ab = x \odot ((y \odot a) \odot b) = x \odot ((y \odot a)b) = x \odot (b(y \odot a)) = x \odot (b \odot (y \odot a)) = (x \odot b) \odot (y \odot a) = (x \odot b)(y \odot a)$. Similarly, we have $(y \odot x)ab = a(y \odot x)b = (a \odot y)(x \odot b) = (x \odot b)(a \odot y) = ((x \odot b)a) \odot y = a((x \odot b) \odot y) = ((x \odot b) \odot y)a = (x \odot b) \odot (y \odot a) = (x \odot b)(y \odot a)$. Hence, $((x \odot y)a)b = ((y \odot x)a)b$ for all $a, b \in S_{\alpha\beta}$. Hence, by the reductivity of $S_{\alpha\beta}$, we have $x \odot y = y \odot x$.

Thus, $S(\circ)$ is commutative. Therefore, $P_r(G)$ is an f.c.e.-property. Next, suppose that there exists an f.c.e.-property $P(G)$ such that $P(G) \not\leq P_r(G)$. Since $P_r(G) \equiv P_r^*(G)$, $P(G) \not\leq P_r^*(G)$. Hence, there is a commutative semigroup S_γ such that $P(S_\gamma)$ is true and $P_r^*(S_\gamma)$ is not true. Since $P_r^*(S_\gamma)$ is not true, there is a system $\{u, v; \xi, \eta\}$ of distinct elements u, v of S_γ and translations ξ, η on S_γ such that $\xi\eta = \eta\xi = \rho_u = \rho_v$, and $\xi(u) = \xi(v)$ and $\eta(u) = \eta(v)$. Let S_α and S_β be infinite cyclic semigroups generated by a and b respectively: $S_\alpha = (a)$ and $S_\beta = (b)$. Let $L_3 = \{\alpha, \beta, \gamma\}$ be a semilattice consisting of three elements α, β, γ such that $\alpha < \gamma, \beta < \gamma, \alpha \not\leq \beta$ and $\beta \not\leq \alpha$. Let $S = S_\alpha \dot{+} S_\beta \dot{+} S_\gamma$. Define multiplication \circ in S as follows: $a^i \circ a^j = a^{i+j}; b^i \circ b^j = b^{i+j}; a \circ b = u, b \circ a = v; x \circ y = xy$ if $x, y \in S_\gamma$; $a^i \circ x = x \circ a^i = \xi^i(x)$ for $x \in S_\gamma$; $b^i \circ x = x \circ b^i = \eta^i(x)$ for $x \in S_\gamma$; and $a^i \circ b^j = \xi^{i-1}\eta^{j-1}(u), b^j \circ a^i = \xi^{i-1}\eta^{j-1}(u)$ if $i+j \geq 3$ (we regard each of ξ^0 and η^0 as the identity mapping on S_γ). Then $S(\circ)$ is a non-commutative composition of $\{S_\alpha, S_\beta, S_\gamma\}$ with respect to L_3 . This contradicts to the assumption that $P(G)$ is an f.c.e.-property and $P(S_\gamma)$ is true. Hence, $P_0(G) \leq P_r(G)$ for all f.c.e.-property $P_0(G)$.

COROLLARY. Let $\Omega = \{S_\xi : \xi \in \chi\}$ (χ : a set) be a collection of commutative semigroups S_ξ , where $P_r(S_\xi)$ is true for each $S_\xi \in \Omega$. Then, every semilattice composition of Ω is commutative.

Proof. Obvious.

Remarks. (1) It is easy to see that if $P_0(G)$ is an f.c.e. [l.c.e.]-property and if $P(G)$ is an abstract property such that $P(G) \leq P_0(G)$, then $P(G)$ is also an f.c.e. [l.c.e.]-property. Let $\mathfrak{P}(Q) = \{P(G) : P(G) \text{ is an abstract property such that } P(G) \leq P_0(G)\}$ and $\mathfrak{P}(R) = \{P(G) : P(G) \text{ is an abstract property such that } P(G) \leq P_r(G)\}$. Then, $\mathfrak{P}(Q)$ and $\mathfrak{P}(R)$ are the set of all l.c.e.-properties and the set of all f.c.e.-properties respectively.

(2) As was shown in Section 1, there exists a universal commutative semigroup S which has a zero element 0 and whose annihilator A contains a non-zero element. Since $P_0(S)$ is true and $P_r(S)$ is not true, $P_0(G) \neq P_r(G)$. Hence, $P_0(G) > P_r(G)$. This also means that quasi-reductivity does not imply reductivity.

(3) Since $P_r(G)$ is weaker than each of separativity (see [2]) and cancellativity, the following results immediately follow from the above-mentioned Corollary:

(i) A semigroup which is a semilattice of commutative reductive semigroups is commutative and reductive.

(ii) A semigroup which is a semilattice of separative commutative semigroups is separative and commutative.

(iii) A semigroup which is a semilattice of cancellative commutative semigroups is separative and commutative.

The converse of the result (iii) also holds (see [2]); i.e., a separative commutative semigroup is a semilattice of cancellative commutative semigroups.

4. MAXIMAL F.C.I. [L.C.I.]-PROPERTIES

In this section, we shall show the existence of a maximal l.c.i.-property and a maximal f.c.i.-property. Especially, it will be proved that $P_q(G)$ is a maximal l.c.i.-property and both $P_r(G)$ and $P_u(G)$ are maximal f.c.i.-properties.

THEOREM 6. *There exist a maximal l.c.i.-property and a maximal f.c.i.-property.*

Proof. Let $\mathcal{F} = \{P_\lambda(G) : \lambda \in \Lambda\}$ be the set of all f.c.i.-properties $P_\lambda(G)$. Then \mathcal{F} is clearly a partially ordered set with respect to the ordering relation \leq defined by (1.6). (Recall that equivalent properties are regarded as the same property). Let $\mathcal{T} = \{P_\tau(G) : \tau \in A_0\}$ be any totally ordered subset of \mathcal{F} . Define an abstract property $T(G)$ as follows: $T(G) = \bigvee_{\tau \in A_0} P_\tau(G)$, i.e., $T(G)$ = the property "being at least one of $\{P_\tau(G) : \tau \in A_0\}$ ". Hence, a commutative semigroup S satisfies $T(G)$ if, and only if, S satisfies at least one of the properties $P_\tau(G)$, $\tau \in A_0$. Now, let $\Omega = \{S_\xi : \xi \in \chi\}$ (χ : a set) be a collection of commutative semigroups S_ξ such that every S_ξ satisfies $T(G)$. Suppose that there exists a non-commutative semilattice composition $S(\circ) = \Sigma\{S_\xi : \xi \in \chi(*)\}$ of Ω . Then there exist a, b such that $a \in S_\gamma$, $b \in S_\delta$, $\gamma, \delta \in \chi$ and $a \circ b \neq b \circ a$. Clearly, both $a \circ b$ and $b \circ a$ are contained in $S_{\gamma * \delta}$. Put $S_\gamma \dot{+} S_\delta \dot{+} S_{\gamma * \delta} = M$. Then $M(\circ)$ is a subsemigroup of $S(\circ)$ and is non-commutative. Since $T(S_\gamma)$, $T(S_\delta)$ and $T(S_{\gamma * \delta})$ are all true, there exist $P_\alpha(G)$, $P_\beta(G)$ and $P_\epsilon(G)$ in the collection $\{P_\tau(G) : \tau \in A_0\}$ such that $P_\alpha(S_\gamma)$, $P_\beta(S_\delta)$ and $P_\epsilon(S_{\gamma * \delta})$ are true. Let $P_\eta(G)$ be the weakest property in $\{P_\alpha(G), P_\beta(G), P_\epsilon(G)\}$. Then $P_\eta(G)$ is of course an f.c.i.-property and $P_\eta(S_\gamma)$, $P_\eta(S_\delta)$, $P_\eta(S_{\gamma * \delta})$ are all true. Hence, the semilattice composition $M(\circ)$ of $\{S_\gamma, S_\delta, S_{\gamma * \delta}\}$ must be commutative. However, this is a contradiction since $M(\circ)$ was non-commutative. Consequently, every semilattice composition of Ω must be commutative. Therefore, $T(G)$ is an f.c.i.-property and hence $T(G) \in \mathcal{F}$. Since $P_\tau(G) \leq T(G)$ for all $\tau \in A_0$, $T(G)$ is an upper bound of \mathcal{T} . Thus, \mathcal{F} is an inductively ordered set. Hence, there exists a maximal f.c.i.-property in \mathcal{F} . The existence of a maximal l.c.i.-property is also proved by a similar method.

COROLLARY. For any f.c.i. [l.c.i.]-property $P(G)$, there exists a maximal f.c.i. [l.c.i.]-property $P_m(G)$ such that $P(G) \leq P_m(G)$.

Proof. This can be proved by an analogous method to Theorem 6.

In fact, the following three theorems show that quasi-reductivity is a maximal l.c.i.-property and both reductivity and universality are maximal f.c.i.-properties:

THEOREM 7. $P_q(G)$ is a maximal l.c.i.-property.

Proof. It is obvious from Remark (1) for Theorem 4 that $P_q(G)$ is an l.c.i.-property. Suppose that there is an l.c.i.-property $P(G)$ such that $P(G) > P_q(G)$. Then, there exists a commutative semigroup S_0 such that $P(S_0)$ is true and $P_q(S_0)$ is not true. Since $P_q(S_0)$ is not true, there is a system $\{\sigma, \rho\}$ of distinct two translations on S_0 such that $\sigma\rho = \rho\sigma$ and $\sigma \mid S_0^2 = \rho \mid S_0^2$. Hence by Lemma 9, there exist distinct two elements x, y and a prime element t in S_0 such that $xa = ya$ for all $a \in S_0$, $\sigma(t) = x$ and $\rho(t) = y$. Let $S_1 = \langle a \rangle$ be an infinite cyclic semigroup generated by a , and let $S_1 \dot{+} S_0 = S$. Define multiplication \circ in S as follows: $a^i \circ u = \rho^i(u)$ for $u \in S_0$; $u \circ a^i = \sigma^i(u)$ for $u \in S_0$; $v \circ w = vw$ if $v, w \in S_1$ or $\in S_0$. Then $S(\circ)$ becomes a non-commutative linear composition of $\{S_1, S_0\}$. Since $P_q(S_1)$ is true and $P_q(G) < P(G)$, $P(S_1)$ is also true. Hence both S_1 and S_0 satisfy $P(G)$, but there is a non-commutative linear composition of $\{S_1, S_0\}$. Thus, we have a contradiction. Hence, there is no l.c.i.-property $P(G)$ such that $P(G) > P_q(G)$.

THEOREM 8. $P_r(G)$ is a maximal f.c.i.-property.

Proof. It is obvious from Corollary to Theorem 5 that $P_r(G)$ is an f.c.i.-property. Suppose that there is an f.c.i.-property $P(G)$ such that $P_r(G) < P(G)$. Then, there exists a commutative semigroup S_γ such that $P(S_\gamma)$ is true and $P_r(S_\gamma)$ is not true. Since $P_r(S_\gamma)$ is not true, there is a system $\{u, v; \xi, \eta\}$ of distinct two elements u, v of S_γ and translations ξ, η on S_γ such that $\xi\eta = \eta\xi = \rho_u = \rho_v$, $\xi(u) = \xi(v)$ and $\eta(u) = \eta(v)$. Let S_α and S_β be infinite cyclic semigroups generated by a and b respectively: $S_\alpha = \langle a \rangle$ and $S_\beta = \langle b \rangle$. Define multiplication \circ in $S = S_\alpha \dot{+} S_\beta \dot{+} S_\gamma$ as follows: $a^i \circ a^j = a^{i+j}$; $b^i \circ b^j = b^{i+j}$; $a \circ b = u$, $b \circ a = v$; $x \circ y = xy$ if $x, y \in S_\gamma$; $a^i \circ x = x \circ a^i = \xi^i(x)$ for $x \in S_\gamma$; $b^i \circ x = x \circ b^i = \eta^i(x)$ for $x \in S_\gamma$; and $a^i \circ b^j = b^j \circ a^i = \xi^{i-1}\eta^{j-1}(u)$ if $i+j \geq 3$ (where ξ^0, η^0 are regarded as the identity mapping on S_γ). Then $S(\circ)$ becomes a non-commutative semilattice composition of $\{S_\alpha, S_\beta, S_\gamma\}$. Since $P_r(S_\alpha)$ and $P_r(S_\beta)$ are true and $P_r(G) < P(G)$, $P(S_\alpha)$ and $P(S_\beta)$ are also true. Hence, the semilattice composition $S(\circ)$ of $\{S_\alpha, S_\beta, S_\gamma\}$ must be commutative since $P(G)$ is an f.c.i.-property

and since each of S_α , S_β and S_γ satisfies $P(G)$. Thus, we have a contradiction. Therefore, there is no f.c.i.-property $P(G)$ such that $P_\nu(G) < P(G)$.

THEOREM 9. $P_u(G)$ is a maximal f.c.i.-property.

Proof. At first, we prove that universality $P_u(G)$ is an f.c.i.-property. Let $\Omega = \{S_\xi : \xi \in \chi\}$ (χ : a set) be any collection of universal commutative semigroups S_ξ . Let $S(\odot) = \mathcal{Z}\{S_\xi : \xi \in \chi(*)\}$ be a semilattice composition of Ω . Let a, b be any two elements of $S = \mathcal{Z}\{S_\xi : \xi \in \chi\}$. Then, there exist $S_\alpha, S_\beta \in \Omega$ such that $S_\alpha \ni a$ and $S_\beta \ni b$. It is obvious that $a \odot b \in S_{\alpha * \beta}$. Since $a \in S_\alpha$ and S_α is universal, $a = a_1 a_2$ for some $a_1, a_2 \in S_\alpha$. Now, clearly each of $a_1 \odot b, b \odot a_1, a_2 \odot b$ and $b \odot a_2$ is contained in $S_{\alpha * \beta}$. $a \odot b = a_1 a_2 \odot b = a_1 \odot (a_2 \odot b) = a_1 \odot xy$ for some $x, y \in S_{\alpha * \beta}$. $a_1 \odot xy = (a_1 \odot x)y = y \odot (a_1 \odot x) = (y \odot a_1)x = x(y \odot a_1) = xy \odot a_1 = (a_2 \odot b) \odot a_1 = a_2 \odot (b \odot a_1) = a_2 \odot (b \odot a_1) = a_2 \odot zw$ for some $z, w \in S_{\alpha * \beta}$. Further, $a_2 \odot zw = (a_2 \odot z)w = w(a_2 \odot z) = (w \odot a_2)z = z(w \odot a_2) = zw \odot a_2 = (b \odot a_1) \odot a_2 = b \odot a_1 a_2 = b \odot a$. Hence, we have $a \odot b = b \odot a$. This means that $S(\odot)$ is commutative. Thus, $P_u(G)$ is an f.c.i.-property. Next, suppose that there is an f.c.i.-property $P(G)$ such that $P_u(G) < P(G)$. Then, there exists a commutative semigroup S_0 such that $P(S_0)$ is true and $P_u(S_0)$ is not true. Let S_γ be a universal commutative semigroup which has a zero element 0 and whose annihilator A contains a nonzero element v . (The existence of such a semigroup S_γ has been shown in Remark of Section 1). Since $P_u(S_\gamma)$ is true and $P_u(G) < P(G)$, $P(S_\gamma)$ is also true. Put $S_\alpha = S_\beta = S_0$, and define multiplication \odot in $S = S_\alpha \dot{+} S_\beta \dot{+} S_\gamma$ as follows: $x \odot y = xy$ if $x, y \in S_\alpha, \in S_\beta$ or $\in S_\gamma$; $S_\alpha \odot S_\gamma = S_\gamma \odot S_\alpha = S_\beta \odot S_\gamma = S_\gamma \odot S_\beta = \{0\}$; $S_\alpha^2 \odot S_\beta = S_\beta \odot S_\alpha^2 = S_\beta^2 \odot S_\alpha = S_\alpha \odot S_\beta^2 = \{0\}$; $S_\alpha^* \odot S_\beta^* = \{v\}$; and $S_\beta^* \odot S_\alpha^* = \{0\}$, where $S_\alpha^* = S_\alpha \setminus S_\alpha^2$, $S_\beta^* = S_\beta \setminus S_\beta^2$ and 0 is the zero element of S_γ . Then, the resulting system $S(\cdot)$ is a non-commutative semilattice composition of Ω . This contradicts to the fact that $P(G)$ is an f.c.i.-property and each of S_α, S_β and S_γ satisfies $P(G)$. Hence, there is no f.c.i.-property $P(G)$ such that $P_u(G) < P(G)$. That is, $P_u(G)$ is a maximal f.c.i.-property.

From Theorem 9, we also have immediately

COROLLARY. *A semigroup which is a semilattice of universal commutative semigroups is universal and commutative.*

Remark. Let $\mathcal{F}[\mathcal{L}]$ be the set of all f.c.i. [l.c.i.]-properties. For $P_1(G), P_2(G) \in \mathcal{F}[\mathcal{L}]$, let us define an abstract property $P_1(G) \wedge P_2(G)$ as follows:

(4.1) $P_1(G) \wedge P_2(G)$ = the property "being both $P_1(G)$ and $P_2(G)$ ". Then, it is easy to see that $P_1(G) \wedge P_2(G) \in \mathcal{F}[\mathcal{L}]$ for any $P_1(G), P_2(G) \in \mathcal{F}[\mathcal{L}]$ and

$P_1(G) \wedge P_2(G)$ is the greatest lower bound of $P_1(G)$ and $P_2(G)$. Further, in fact $\mathcal{F}[\mathcal{L}]$ is a semilattice with respect to this operation \wedge .

Since $P_u(G)$ and $P_r(G)$ are non-equivalent maximal f.c.i.-properties, it is obvious that there is no greatest f.c.i.-property, i.e., there is no weakest f.c.i.-property $P_g(G)$ in the following sense:

$$(4.2) \quad P(G) \leq P_g(G) \text{ for any f.c.i.-property } P(G).$$

However, the authors can not solve the following two problems and leave them as open problems:

Problem 1. Is there a maximal l.c.i.-property except $P_q(G)$? That is: Is $P_q(G)$ the greatest (weakest) l.c.i.-property? Determine all of the maximal l.c.i.-properties.

Problem 2. Is there a maximal f.c.i.-property except $P_u(G)$ and $P_r(G)$? Determine all of the maximal f.c.i.-properties.

As a partial solution of Problem 1, we obtain the following result: Let C be an infinite cyclic semigroup: $C = \{a, a^2, \dots, a^n, \dots\}$. Let C^1 be the adjunction of an identity element to C : $C^1 = C \dot{+} \{1\}$. Let \mathcal{L}_0 be the set of all l.c.i.-properties $L(G)$ satisfied by C^1 , and \mathcal{L}^* the set of all abstract properties $P(G)$ such that $P(G) \leq L(G)$ for some $L(G) \in \mathcal{L}_0$. Then $\mathcal{L}^* \ni P_q(G), P_r(G), P_u(G)$, since each of $P_q(C^1)$, $P_u(C^1)$ and $P_r(C^1)$ is true and each of $P_q(G)$, $P_u(G)$ and $P_r(G)$ is an l.c.i.-property. Further, $\mathcal{L}^* \supset \mathcal{L} = \{P(G) : P(G) \text{ is an l.c.i.-property which is comparable with } P_r(G) \text{ or } P_u(G)\}$. In fact, let $P(G)$ be a property contained in \mathcal{L} . If $P(G) \leq P_r(G)$ or $\leq P_u(G)$, then $P(G) \in \mathcal{L}^*$ since each of $P_r(G)$ and $P_u(G)$ is an l.c.i.-property and is satisfied by C^1 . If $P(G) > P_r(G)$ or $> P_u(G)$, then $P(C^1)$ is true since each of $P_r(C^1)$ and $P_u(C^1)$ is true. Since $P(C^1)$ is true and $P(G)$ is an l.c.i.-property, $P(G)$ is also contained in \mathcal{L}^* . In any case, $P(G) \in \mathcal{L}^*$. Therefore, $\mathcal{L} \subset \mathcal{L}^*$. Especially, cancellativity, separativity, regularity and the property "being a commutative semigroup G with 1" are all contained in \mathcal{L}^* .

Now, we have

THEOREM 10. $P_q(G)$ is the greatest (i.e. weakest) l.c.i.-property in \mathcal{L}^* .

Proof. Let $P(G)$ be an l.c.i.-property contained in \mathcal{L}^* . Then, there exists an l.c.i.-property $L(G)$ such that $L(C^1)$ is true and $P(G) \leq L(G)$. Suppose that $L(G) \not\leq P_q(G)$. Then, there exists a commutative semigroup S_0 such that $L(S_0)$ is true and $P_q(S_0)$ is not true. Since $P_q(S_0)$ is not true, there exists a system $\{\sigma, \rho\}$ of distinct two translations σ, ρ on S_0 such that $\sigma\rho = \rho\sigma$ and $\sigma \mid S_0^2 = \rho \mid S_0^2$. Let $S_1 = C$ (= the infinite cyclic semigroup (a)), and define multiplication \circ in $S^* = S_1 \dot{+} S_0$ as follows: $a^i \circ u = \rho^i(u)$ for $u \in S_0$; $u \circ a^i = \sigma^i(u)$ for $u \in S_0$; and $v \circ w = vw$ if $v, w \in S_1$ or $\in S_0$.

Then, $S^*(\circ)$ is a non-commutative linear composition of $\{S_1, S_0\}$. Let S be the adjunction of an identity element to $S^*(\circ) : S = S^* \dot{+} \{1\} = C^1 \dot{+} S_0$. Then S is clearly a non-commutative linear composition of $\{C^1, S_0\}$. This contradicts to our assumption that $L(C^1)$, $L(S_0)$ are true and $L(G)$ is an l.c.i.-property. Hence $L(G) \leq P_q(G)$, and accordingly $P(G) \leq L(G) \leq P_q(G)$. This means that $P_q(G)$ is the greatest l.c.i.-property in \mathcal{L}^* .

ACKNOWLEDGMENTS

The authors are indebted to Professor Takayuki Tamura for valuable discussions and for substantial help in the development of the material of this paper.

REFERENCES

1. CLIFFORD, A. H. Naturally totally ordered commutative semigroups. *Amer. J. Math.* **126** (1954), 631–646.
2. CLIFFORD, A. H., AND PRESTON, G. B. "The Algebraic Theory of Semigroups." Vol. 1 [Mathematical Surveys of the American Mathematical Society (7th in Series)]. American Mathematical Society, Providence, Rhode Island, 1961.
3. COHN, P. M. "Universal Algebra." Harper & Row, New York, Evanston, and London, 1965.
4. YAMADA, M. Compositions of semigroups. *Kōdai Math. Sem. Rep.* **8** (1956), 107–111.
5. YOSHIDA, R. Ideal extensions of semigroups and compound semigroups. *Mem. Res. Inst. Sci. Engrg., Ritumeikan Univ.* **13** (1965), 1–8.